

Dental Quiz for General Practitioners

Compiled by

Dr. Ashmey Mariya Sajan¹, Dr. Hima John²

¹House surgeon 2013-14 batch,
Annoor Dental College & Hospital, Muvattupuzha.

²Senior lecturer, Dept of Oral Medicine and Radiology,
Annoor Dental College & Hospital, Muvattupuzha.

- The tissue of which lesion has been described as resembling a ' blood soaked sponge with large pores '?
 - Cavernous hemangioma
 - Capillary hemangioma
 - Aneurysmal bone cyst
 - Eruption bone cyst
- The radiographic appearance of Botryoid odontogenic cyst is
 - Apple green appearance
 - Grape bunch appearance
 - Lemon peel appearance
 - Orange peel appearance
- In case of conscious sedation or children aged 6 months to 3 years clear liquids should be stopped. (OR) According to AAPD guidelines , for a child undergoing sedation procedure, the human breast milk should be stopped at least _____ hours before the procedure .
 - 4 hours , before the procedure
 - 6 hours , before the procedure
 - 8 hours , before the procedure
 - 10 hours , before the procedure
- Most common unfavourable sequelae of serial extraction
 - Disto-axial inclination of canine
 - Residual space at the extraction site
 - Mesio - axial inclination of second premolar
 - Increase in overbite
- Which of the following drug is an antipseudomonal penicillin ?
 - Cephalexin
 - Cloxacillin
 - Piperacillin
 - Dicloxacillin
- The diameter of the tip of periodontal probe is :
 - 0.25mm
 - 0.75mm
 - 0.5mm
 - 1mm
- The dominant histologic feature of infarction is
 - Liquefactive necrosis
 - Coagulative necrosis
 - Chronic inflammation
 - Scar tissue
- Vitremer is
 - Resin modified GIC
 - Metal modified GIC
 - Dentin bonding agent
 - All of the above

9. Broca's area is concerned with
 - a) Word formation
 - b) Comprehends language not speaking
 - c) Present in posterior part of temporal lobe
 - d) Is the confluence of somatic , auditory and visceral areas
10. The following is the first sign appeared in shock
 - a) Hypotension
 - b) Tachycardia
 - c) Hypertension
 - d) Syncope
11. First local anaesthesia to be used clinically was
 - a) Cocaine
 - b) Bupivacaine
 - c) Procaine
 - d) Lignocaine
12. Reversal lines which may be seen on the cribriform plate (alveolar bone proper) of the alveolar process indicate the cessation of
 - a) Osteoblastic activity
 - b) Osteoclastic activity
 - c) Myeloid activity
 - d) Healing activity
13. The porcelain constituent that fuses at firing temperature and forms a matrix is
 - a) Kaolin
 - b) Quartz
 - c) Feldspar
 - d) Aluminium oxide
14. Krebs' cycle occurs in _____ conditions.
 - a) Aerobic
 - b) Anaerobic
 - c) Microaerophilic
 - d) Aerobic and anaerobic
15. Phosphofructokinase is the key enzyme of :
 - a) Glycolysis
 - b) Gluconeogenesis
 - c) Beta oxidation
 - d) TCA cycle
16. The serum concentration of which of the following human IgG subclass is maximum?
 - a) IgG1
 - b) IgG2
 - c) IgG3
 - d) IgG4
17. Which of the following methods is used to sterilize gutta-percha?
 - a) Gas sterilization
 - b) Steam sterilization
 - c) Ethyl alcohol
 - d) Quarternary ammonium compounds
18. Pulp capillary pressure is:
 - a) 14cm of H₂O
 - b) 25cm of H₂O
 - c) 80 cm of H₂O
 - d) 120 cm of H₂O
19. Crude mortality rate is :
 - a) Per 100
 - b) Per 1000
 - c) Per 10000
 - d) Per 100000
20. The first costochondral joint is a:
 - a) Fibrous joint
 - b) Synovial joint
 - c) Synarthrosis
 - d) Ball and socket joint

ANSWERS:

1.c) , 2.b) , 3.a) , 4.d) , 5.c) , 6.c) , 7.b) , 8.a) , 9.a) , 10.b) , 11.a) , 12.b) , 13.c) , 14.a) , 15.a) , 16.a) , 17.c) , 18.c) , 19.b) , 20.c).